



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer



**Measurement of the Difference of Time-Integrated CPAsymmetries in  $D^0KK^+$  and  $D^0+$  Decays**



# Measurement of the Difference of Time-Integrated $CP$ Asymmetries in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ Decays

R. Aaij *et al.*\*

(LHCb Collaboration)

(Received 11 February 2016; published 9 May 2016)

A search for  $CP$  violation in  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  decays is performed using  $pp$  collision data, corresponding to an integrated luminosity of  $3 \text{ fb}^{-1}$ , collected using the LHCb detector at center-of-mass energies of 7 and 8 TeV. The flavor of the charm meson is inferred from the charge of the pion in  $D^{*+} \rightarrow D^0 \pi^+$  and  $D^{*-} \rightarrow \bar{D}^0 \pi^-$  decays. The difference between the  $CP$  asymmetries in  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  decays,  $\Delta A_{CP} \equiv A_{CP}(K^- K^+) - A_{CP}(\pi^- \pi^+)$ , is measured to be  $[-0.10 \pm 0.08(\text{stat}) \pm 0.03(\text{syst})]\%$ . This is the most precise measurement of a time-integrated  $CP$  asymmetry in the charm sector from a single experiment.

DOI: 10.1103/PhysRevLett.116.191601

Violation of charge-parity ( $CP$ ) symmetry in weak decays of hadrons is described in the Standard Model (SM) by the Cabibbo-Kobayashi-Maskawa (CKM) matrix and has been observed in  $K$ - and  $B$ -meson systems [1–5]. However, no  $CP$  violation has been observed in the charm sector, despite the experimental progress seen in charm physics in the last decade. Examples are the unambiguous observation of  $D^0$ – $\bar{D}^0$  meson mixing [6–11], and measurements of  $CP$  asymmetry observables in  $D$  meson decays, reaching an experimental precision of  $\mathcal{O}(10^{-3})$  [12]. The amount of  $CP$  violation is expected to be below the percent level [13–20], but large theoretical uncertainties due to long distance interactions prevent precise SM calculations. Charm hadrons provide a unique opportunity to search for  $CP$  violation with particles containing only up-type quarks.

This Letter presents a measurement of the difference between the time-integrated  $CP$  asymmetries of  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  decays, performed with  $pp$  collision data corresponding to an integrated luminosity of  $3 \text{ fb}^{-1}$  collected using the LHCb detector at center-of-mass energies of 7 and 8 TeV. The inclusion of charge-conjugate decay modes is implied throughout except in the definition of asymmetries. This result is an update of the previous LHCb measurement with  $0.6 \text{ fb}^{-1}$  of data, in which a value of  $\Delta A_{CP} = (-0.82 \pm 0.21)\%$  was obtained [21].

The time-dependent  $CP$  asymmetry,  $A_{CP}(f; t)$ , for  $D^0$  mesons decaying to a  $CP$  eigenstate  $f$  is defined as

$$A_{CP}(f; t) \equiv \frac{\Gamma(D^0(t) \rightarrow f) - \Gamma(\bar{D}^0(t) \rightarrow f)}{\Gamma(D^0(t) \rightarrow f) + \Gamma(\bar{D}^0(t) \rightarrow f)}, \quad (1)$$

where  $\Gamma$  denotes the decay rate. For  $f = K^- K^+$  and  $f = \pi^- \pi^+$ ,  $A_{CP}(f; t)$  can be expressed in terms of a direct component associated with  $CP$  violation in the decay amplitudes, and an indirect component associated with  $CP$  violation in the mixing or in the interference between mixing and decay. In the limit of exact symmetry under a transformation interchanging  $d$  and  $s$  quarks ( $U$ -spin symmetry), the direct component is expected to be equal in magnitude and opposite in sign for  $K^- K^+$  and  $\pi^- \pi^+$  decays [22]. However, large  $U$ -spin breaking effects could be present [13,16,23,24].

The measured time-integrated asymmetry,  $A_{CP}(f)$ , depends upon the reconstruction efficiency as a function of the decay time. It can be written as [25,26]

$$A_{CP}(f) \approx a_{CP}^{\text{dir}}(f) \left( 1 + \frac{\langle t(f) \rangle}{\tau} y_{CP} \right) + \frac{\langle t(f) \rangle}{\tau} a_{CP}^{\text{ind}}, \quad (2)$$

where  $\langle t(f) \rangle$  denotes the mean decay time of  $D^0 \rightarrow f$  decays in the reconstructed sample,  $a_{CP}^{\text{dir}}(f)$  as the direct  $CP$  asymmetry,  $\tau$  the  $D^0$  lifetime,  $a_{CP}^{\text{ind}}$  the indirect  $CP$  asymmetry, and  $y_{CP}$  is the deviation from unity of the ratio of the effective lifetimes of decays to flavor specific and  $CP$ -even final states. To a good approximation,  $a_{CP}^{\text{ind}}$  is independent of the decay mode [22,27].

Neglecting terms of the order  $\mathcal{O}(10^{-6})$ , the difference in  $CP$  asymmetries between  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  is

$$\begin{aligned} \Delta A_{CP} &\equiv A_{CP}(K^- K^+) - A_{CP}(\pi^- \pi^+) \\ &\approx \Delta a_{CP}^{\text{dir}} \left( 1 + \frac{\langle \bar{t} \rangle}{\tau} y_{CP} \right) + \frac{\Delta \langle t \rangle}{\tau} a_{CP}^{\text{ind}}, \end{aligned} \quad (3)$$

\*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the Creative Commons Attribution 3.0 License. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

where  $\overline{\langle t \rangle}$  is the arithmetic average of  $\langle t(K^-K^+) \rangle$  and  $\langle t(\pi^-\pi^+) \rangle$ .

The most precise measurements of the time-integrated  $CP$  asymmetries in  $D^0 \rightarrow K^-K^+$  and  $D^0 \rightarrow \pi^-\pi^+$  decays to date have been performed by the LHCb [21,28], CDF [29], BABAR [30] and Belle [31,32] collaborations. The measurement in Ref. [28] uses  $D^0$  mesons produced in semileptonic  $b$ -hadron decays, where the charge of the muon is used to identify the flavor of the  $D^0$  meson at production, while the other measurements use  $D^0$  mesons produced in the decay of the  $D^*(2010)^+$  meson, hereafter referred to as  $D^{*+}$ .

The raw asymmetry,  $A_{\text{raw}}(f)$ , measured for  $D^0$  decays to a final state  $f$  is defined as

$$A_{\text{raw}}(f) \equiv \frac{N(D^{*+} \rightarrow D^0(f)\pi_s^+) - N(D^{*-} \rightarrow \bar{D}^0(f)\pi_s^-)}{N(D^{*+} \rightarrow D^0(f)\pi_s^+) + N(D^{*-} \rightarrow \bar{D}^0(f)\pi_s^-)}, \quad (4)$$

where  $N$  is the number of reconstructed signal candidates of the given decay and the flavor of the  $D^0$  meson is identified using the charge of the soft pion ( $\pi_s^+$ ) in the strong decay  $D^{*+} \rightarrow D^0\pi_s^+$ . The raw asymmetry can be written, up to  $\mathcal{O}(10^{-6})$ , as

$$A_{\text{raw}}(f) \approx A_{CP}(f) + A_D(f) + A_D(\pi_s^+) + A_P(D^{*+}), \quad (5)$$

where  $A_D(f)$  and  $A_D(\pi_s^+)$  are the asymmetries in the reconstruction efficiencies of the  $D^0$  final state and of the soft pion, and  $A_P(D^{*+})$  is the production asymmetry for  $D^{*+}$  mesons, arising from the hadronization of charm quarks in  $pp$  collisions. The magnitudes of  $A_P(D^{*+})$  [33] and  $A_D(\pi_s^+)$  [34] are both about 1%. Equation (5) is only valid when reconstruction efficiencies of the final state  $f$  and of the soft pion are independent. Since both  $K^-K^+$  and  $\pi^-\pi^+$  final states are self-conjugate,  $A_D(K^-K^+)$  and  $A_D(\pi^-\pi^+)$  are identically zero. To a good approximation  $A_D(\pi_s^+)$  and  $A_P(D^{*+})$  are independent of the final state  $f$  in any given kinematic region, and thus cancel in the difference, giving

$$\Delta A_{CP} = A_{\text{raw}}(K^-K^+) - A_{\text{raw}}(\pi^-\pi^+). \quad (6)$$

However, to take into account an imperfect cancellation of detection and production asymmetries due to the difference in the kinematic properties of the two decay modes, the kinematic distributions of  $D^{*+}$  mesons decaying to the  $K^-K^+$  final state are reweighted to match those of  $D^{*+}$  mesons decaying to the  $\pi^-\pi^+$  final state. The weights are calculated for each event using the ratios of the background-subtracted distributions of the  $D^{*+}$  momentum, transverse momentum, and azimuthal angle for both final states after the final selection.

The LHCb detector [35,36] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ ,

designed for the study of particles containing  $b$  or  $c$  quarks. The two ring-imaging Cherenkov detectors [37] provide particle identification (PID) to distinguish kaons from pions for momenta ranging from a few GeV/ $c$  to about 100 GeV/ $c$ . The direction of the field polarity (up or down) of the LHCb dipole magnet is reversed periodically, giving data samples of comparable size for both magnet polarities.

To select  $D^{*+}$  candidates, events must satisfy hardware and software trigger requirements and a subsequent offline selection. The trigger consists of a hardware stage, based on high transverse momentum signatures in the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. When the hardware trigger decision is initiated by calorimeter deposits from  $D^0$  decay products, the event is categorized as “triggered on signal” (TOS). Events that are not TOS, but in which the hardware trigger decision is due to particles in the event other than the  $D^{*+}$  decay products, are also accepted; these are referred to as “not triggered on signal” (nTOS). The events associated with these trigger categories present different kinematic properties. To have cancellation of production and detection asymmetries the data are split into TOS and nTOS samples and  $\Delta A_{CP}$  is measured separately in each sample.

Both the software trigger and subsequent event selection use kinematic variables and decay time to isolate the signal decays from the background. Candidate  $D^0$  mesons must have a decay vertex that is well separated from all primary  $pp$  interaction vertices (PVs). They are combined with pion candidates to form  $D^{*+}$  candidates. Requirements are placed on the track fit quality, the  $D^{*+}$  vertex fit quality, where the vertex formed by  $D^0$  and  $\pi_s^+$  candidates is constrained to coincide with the associated PV [38], the  $D^0$  transverse momentum and its decay distance, the angle between the  $D^0$  momentum in the laboratory frame and the momentum of the kaon or the pion in the  $D^0$  rest frame, and the smallest impact parameter chi-squared (IP  $\chi^2$ ) of both the  $D^0$  candidate and its decay products with respect to all PVs in the event. The IP  $\chi^2$  is defined as the difference between the  $\chi^2$  of the PV reconstructed with and without the considered particle. Cross-feed backgrounds from  $D$  meson decays with a kaon misidentified as a pion, and vice versa, are reduced using PID requirements. After these selection criteria, the dominant background consists of genuine  $D^0$  candidates paired with unrelated pions originating from the interaction vertex.

Fiducial requirements are imposed to exclude kinematic regions having a large asymmetry in the soft pion reconstruction efficiency (see Figs. 1 and 2 in Ref. [39]). These regions occur because low momentum particles of one charge at large (small) angles in the horizontal plane may be deflected out of the detector acceptance (into the noninstrumented beam pipe region) whereas particles with the other charge are more likely to remain within the




FIG. 1. Fit to the  $\delta m$  spectra, where the  $D^0$  is reconstructed in the final state (left)  $K^-K^+$  and (right)  $\pi^-\pi^+$ . The dashed line corresponds to the background component in the fit.

acceptance. About 70% of the selected candidates are retained by these fiducial requirements.

The candidates satisfying the selection criteria are accepted for further analysis if the mass difference  $\delta m \equiv m(h^+h^-\pi_s^+) - m(h^+h^-) - m(\pi^+)$  for  $h = K, \pi$  is in the range  $0.2 - 12.0 \text{ MeV}/c^2$  and the invariant mass of the  $D^0$  candidate is within 2 standard deviations from the central value of the mass resolution model. The standard deviation corresponds to about  $8 \text{ MeV}/c^2$  and  $10 \text{ MeV}/c^2$  for  $D^0 \rightarrow K^-K^+$  and  $D^0 \rightarrow \pi^-\pi^+$  decays, respectively.

The data sample includes events with multiple  $D^{*+}$  candidates. The majority of these events contain the same reconstructed  $D^0$  meson combined with different soft pion candidates. The fraction of events with multiple candidates in a range of  $\delta m$  corresponding to  $4.0 - 7.5 \text{ MeV}/c^2$  is about 1.2% for TOS events and 2.4% for nTOS events; these fractions are the same for the  $K^-K^+$  and  $\pi^-\pi^+$  final states, and for both magnet polarities. The events with multiple candidates are retained and a systematic uncertainty is assessed.

Signal yields and  $A_{\text{raw}}(K^-K^+)$  and  $A_{\text{raw}}(\pi^-\pi^+)$  are obtained from minimum  $\chi^2$  fits to the binned  $\delta m$  distributions of the  $D^0 \rightarrow K^-K^+$  and  $D^0 \rightarrow \pi^-\pi^+$  samples. The data samples are split into eight mutually exclusive subsamples separated by center-of-mass energy, magnet polarity, and trigger category. The signal shape is studied using simulated data and described by the sum of




FIG. 2. Contour plot of  $\Delta a_{CP}^{\text{dir}}$  versus  $a_{CP}^{\text{ind}}$ . The point at  $(0,0)$  denotes the hypothesis of no  $CP$  violation. The solid bands represent the measurements in Refs. [28,45,46] and the one reported in this Letter. The value of  $y_{CP}$  is taken from Ref. [47]. The contour lines show the 68%, 95%, and 99% confidence-level intervals from the combination.

two Gaussian functions with a common mean, and a Johnson  $S_U$  function [40]. The background is described by an empirical function of the form  $1 - \exp[(\delta m - \delta m_0)/\alpha] + \beta(\delta m/\delta m_0 - 1)$ , where  $\delta m_0$  controls the threshold of the function, and  $\alpha$  and  $\beta$  describe its shape. The fits to the eight subsamples and between the  $K^-K^+$  and  $\pi^-\pi^+$  final states are independent. Fits to the  $\delta m$  distributions corresponding to the whole data sample are shown in Fig. 1.

The  $D^{*+}$  signal yield is  $7.7 \times 10^6$  for  $D^0 \rightarrow K^-K^+$  decays, and  $2.5 \times 10^6$  for  $D^0 \rightarrow \pi^-\pi^+$  decays. The signal purity is  $(88.7 \pm 0.1)\%$  for  $D^0 \rightarrow K^-K^+$  candidates, and  $(87.9 \pm 0.1)\%$  for  $D^0 \rightarrow \pi^-\pi^+$  candidates, in a range of  $\delta m$  corresponding to  $4.0 - 7.5 \text{ MeV}/c^2$ . The fits do not distinguish between the signal and the backgrounds that peak in  $\delta m$ . Such backgrounds, which can arise from  $D^{*+}$  decays where the correct soft pion is found but the  $D^0$  meson is misreconstructed, are suppressed by the PID requirements to less than 4% of the number of signal events in the case of  $D^0 \rightarrow K^-K^+$  decays and to a negligible level in the case of  $D^0 \rightarrow \pi^-\pi^+$  decays. Examples of such backgrounds are  $D^{*+} \rightarrow D^0(K^-\pi^+\pi^0)\pi_s^+$  and  $D^{*+} \rightarrow D^0(\pi^-e^+\nu_e)\pi_s^+$  decays. The effect on  $\Delta A_{CP}$  of residual peaking backgrounds is evaluated as a systematic uncertainty.

The value of  $\Delta A_{CP}$  is determined in each subsample (see Table 1 in Ref. [39]). Testing the eight independent measurements for mutual consistency gives  $\chi^2/\text{ndf} = 6.2/7$ , corresponding to a  $p$ -value of 0.52. The weighted average of the values corresponding to all subsamples is calculated as  $\Delta A_{CP} = (-0.10 \pm 0.08)\%$ , where the uncertainty is statistical.



The central value is considerably closer to zero than  $\Delta A_{CP} = (-0.82 \pm 0.21)\%$ , obtained in our previous analysis where a data sample corresponding to an integrated luminosity of  $0.6 \text{ fb}^{-1}$  was considered [21]. Several factors contribute to the change, including the increased size of the data sample and changes in the detector calibration and reconstruction software. To estimate the impact of processing data using different reconstruction software, the data used in Ref. [21] are divided into three samples. The first (second) sample contains events that are selected when using the old (new) version of the reconstruction software and are discarded by the new (old) one, while the third sample consists of those events that are selected by both versions. The measured values are  $\Delta A_{CP} = (-1.10 \pm 0.46)\%$ ,  $\Delta A_{CP} = (0.13 \pm 0.37)\%$ , and  $\Delta A_{CP} = (-0.71 \pm 0.26)\%$ , respectively. The measurement obtained using the additional data based on an integrated luminosity of  $2.4 \text{ fb}^{-1}$  corresponds to a value of  $\Delta A_{CP} = (-0.06 \pm 0.09)\%$ . A comparison of the four independent measurements gives  $\chi^2/\text{ndf} = 10.5/3$ , equivalent to a  $p$ -value of 0.015. Although this value is small, no evidence of incompatibility among the various subsamples has been found. Only statistical uncertainties are considered in this study.

Many sources of systematic uncertainty that may affect the determination of  $\Delta A_{CP}$  are considered. The possibility of an incorrect description of the signal mass model is investigated by replacing the function in the baseline fit with alternative models that provide equally good descriptions of the data. A value of 0.016% is assigned as systematic uncertainty, corresponding to the largest variation observed using the alternative functions.

To evaluate the systematic uncertainty related to the presence of multiple candidates in an event,  $\Delta A_{CP}$  is measured in samples where one candidate per event is randomly selected. This procedure is repeated 100 times with a different random selection. The difference of the mean value of these measurements from the nominal result, 0.015%, is taken as systematic uncertainty.

A systematic uncertainty associated with the presence of background peaking in the  $\delta m$  signal distribution and not in the  $D^0$  invariant mass distribution is determined by measuring  $\Delta A_{CP}$  from fits to the  $D^0$  invariant mass spectra instead of  $\delta m$ . Fits are made for  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  candidates within a  $\delta m$  window  $4.0\text{--}7.5 \text{ MeV}/c^2$ . The background due to genuine  $D^0$  mesons paired with unrelated pions originating from the interaction vertex is subtracted by means of analogous fits to the candidates in the  $\delta m$  window  $8.0\text{--}12.0 \text{ MeV}/c^2$ , where the signal is not present. The difference in the  $\Delta A_{CP}$  value from the baseline, 0.011%, is assigned as a systematic uncertainty. A systematic uncertainty of 0.004% is assigned for uncertainties associated with the weights calculated for the kinematic reweighting procedure.

A systematic uncertainty is associated with the choice of fiducial requirements on the soft pion applied to exclude

regions with large raw asymmetries. To evaluate this uncertainty, the baseline results are compared to results obtained when looser fiducial requirements are applied. The resulting samples include events closer to the regions with large raw asymmetries, at the edges of the detector acceptance and around the beam pipe (see Fig. 1 in Ref. [39]). The difference in the  $\Delta A_{CP}$  values, 0.017%, is taken as the systematic uncertainty.

Although suppressed by the requirement that the  $D^0$  trajectory points back to the primary vertex,  $D^{*+}$  mesons produced in the decays of beauty hadrons (secondary charm decays) are still present in the final sample. As the  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  decays may have different amounts of this contamination, the value of  $\Delta A_{CP}$  may be biased because of an incomplete cancellation of the production asymmetries of beauty and charm hadrons. The fractions of secondary charm decays are estimated by performing a fit to the distribution of IP  $\chi^2$  of the  $D^0$  with respect to all PVs in the event, and are found to be  $(2.8 \pm 0.1)\%$  and  $(3.4 \pm 0.1)\%$  for the  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  samples, respectively. Using the LHCb measurements of production asymmetries [33,41–43], the corresponding systematic uncertainty is estimated to be 0.004%.

To investigate other sources of systematic uncertainty, numerous robustness checks have been made. The value of  $\Delta A_{CP}$  is studied as a function of data taking periods and no evidence of any dependence is found. A measurement of  $\Delta A_{CP}$  using more restrictive PID requirements is performed, and all variations of  $\Delta A_{CP}$  are found to be compatible within statistical uncertainties. To check for possible reconstruction biases, the stability of  $\Delta A_{CP}$  is also investigated as a function of many reconstructed quantities, including the number of reconstructed PVs, the  $D^0$  invariant mass, the  $D^0$  transverse momentum, the  $D^0$  flight distance, the  $D^0$  azimuthal angle, the smallest IP  $\chi^2$  impact parameter of the  $D^0$  and of the soft pion with respect to all the PVs in the events, the quality of  $D^{*+}$  vertex, the transverse momentum of the soft pion, and the quantity  $\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2}$ , where  $\Delta\phi$  and  $\Delta\eta$  are the differences between  $D^0$  and soft pion azimuthal angles and pseudorapidities. No evidence of dependence of  $\Delta A_{CP}$  on any of these variables is found. An additional cross-check concerns the measured value of  $\Delta A_{\text{bkg}}$ , defined as the difference between the background raw asymmetries  $A_{\text{bkg}}(K^- K^+)$  and  $A_{\text{bkg}}(\pi^- \pi^+)$ . A value of  $\Delta A_{\text{bkg}} = (-0.46 \pm 0.13)\%$  is obtained from the fits. In the absence of misidentified or misreconstructed backgrounds, one would expect a value consistent with zero. Decays of  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  have different sources of backgrounds that do not peak in  $\delta m$ . These include three-body decays of charmed hadrons with misidentified particles in the final state, as well as four-body decays where one particle is not reconstructed. More restrictive PID requirements have been applied to suppress such

backgrounds, and the region of the fits has been extended up to  $16 \text{ MeV}/c^2$  to improve the precision. A value of  $\Delta A_{\text{bkg}} = (-0.22 \pm 0.13)\%$  is found. The corresponding  $\Delta A_{CP}$  value is  $(-0.12 \pm 0.09)\%$ , consistent with the baseline result when the overlap of the two samples is taken into account. Hence, the measurement of  $\Delta A_{CP}$  is robust and is not influenced by the background asymmetry. All contributions are summed in quadrature to give a total systematic uncertainty of  $0.03\%$ .

To interpret the  $\Delta A_{CP}$  result in terms of direct and indirect  $CP$  violation, the reconstructed decay time averages, for  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  samples, are measured. The difference and the average of the mean decay times relative to the  $D^0$  lifetime are computed, giving  $\Delta\langle t \rangle/\tau(D^0) = 0.1153 \pm 0.0007(\text{stat}) \pm 0.0018(\text{syst})$  and  $\langle t \rangle/\tau(D^0) = 2.0949 \pm 0.0004(\text{stat}) \pm 0.0159(\text{syst})$ . The systematic uncertainties are due to the uncertainty on the world average of the  $D^0$  lifetime [44], decay-time resolution model, and the presence of secondary  $D^0$  mesons from  $b$ -hadron decays. Given the dependence of  $\Delta A_{CP}$  on the direct and indirect  $CP$  asymmetries [Eq. (3)] and the measured value of  $\Delta\langle t \rangle/\tau$ , the contribution from indirect  $CP$  violation is suppressed and  $\Delta A_{CP}$  is primarily sensitive to direct  $CP$  violation. Assuming that indirect  $CP$  violation is independent of the  $D^0$  final state, and combining the measurement reported in this Letter with those reported in Ref. [28] and with the LHCb measurements of indirect  $CP$  asymmetries ( $A_\Gamma \simeq -a_{CP}^{\text{ind}}$ ) [45,46] and  $y_{CP}$  [47], the values of the direct and indirect  $CP$  asymmetries are found to be  $a_{CP}^{\text{ind}} = (0.058 \pm 0.044)\%$  and  $\Delta a_{CP}^{\text{dir}} = (-0.061 \pm 0.076)\%$ . Results are summarized in the  $(\Delta a_{CP}^{\text{dir}}, a_{CP}^{\text{ind}})$  plane shown in Fig. 2. The result is consistent with the hypothesis of  $CP$  symmetry with a  $p$ -value of  $0.32$ .

In summary, the difference of time-integrated  $CP$  asymmetries between  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  decays is measured using  $pp$  collision data corresponding to an integrated luminosity of  $3.0 \text{ fb}^{-1}$ . The final result is

$$\Delta A_{CP} = [-0.10 \pm 0.08(\text{stat}) \pm 0.03(\text{syst})]\%,$$

which supersedes the previous result obtained using the same decay channels based on an integrated luminosity of  $0.6 \text{ fb}^{-1}$  [21]. This is the most precise measurement of a time-integrated  $CP$  asymmetry in the charm sector from a single experiment.

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); FOM and NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinECo (Spain); SNSF and SER (Switzerland);

NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), and The Royal Society, Royal Commission for the Exhibition of 1851 and the Leverhulme Trust (United Kingdom).

- 
- [1] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, Evidence for the  $2\pi$  Decay of the  $K_2^0$  Meson, *Phys. Rev. Lett.* **13**, 138 (1964).
  - [2] B. Aubert *et al.* (BABAR Collaboration), Observation of Direct  $CP$  Violation in  $B^0 \rightarrow K^+ \pi^-$  Decays, *Phys. Rev. Lett.* **93**, 131801 (2004).
  - [3] Y. Chao *et al.* (Belle Collaboration), Evidence for Direct  $CP$  Violation in  $B^0 \rightarrow K^+ \pi^-$  Decays, *Phys. Rev. Lett.* **93**, 191802 (2004).
  - [4] R. Aaij *et al.* (LHCb Collaboration), First Observation of  $CP$  Violation in the Decays of  $B_s^0$  Mesons, *Phys. Rev. Lett.* **110**, 221601 (2013).
  - [5] R. Aaij *et al.* (LHCb Collaboration), Observation of  $CP$  violation in  $B^\pm \rightarrow DK^\pm$  decays, *Phys. Lett. B* **712**, 203 (2012); **713**, 351(E) (2012).
  - [6] B. Aubert *et al.* (BABAR Collaboration), Evidence for  $D^0$ - $\bar{D}^0$  Mixing, *Phys. Rev. Lett.* **98**, 211802 (2007).
  - [7] M. Staric *et al.* (Belle Collaboration), Measurement of  $D^0$ - $\bar{D}^0$  mixing and search for  $CP$  violation in  $D^0 \rightarrow K^+ K^-, \pi^+ \pi^-$  decays with the full Belle data set, *Phys. Lett. B* **753**, 412 (2016).
  - [8] T. Aaltonen *et al.* (CDF Collaboration), Evidence for  $D^0$ - $\bar{D}^0$  Mixing Using the CDF II Detector, *Phys. Rev. Lett.* **100**, 121802 (2008).
  - [9] R. Aaij *et al.* (LHCb Collaboration), Observation of  $D^0$ - $\bar{D}^0$  Oscillations, *Phys. Rev. Lett.* **110**, 101802 (2013).
  - [10] R. Aaij *et al.* (LHCb Collaboration), Measurement of  $D^0$ - $\bar{D}^0$  Mixing Parameters and Search for  $CP$  Violation Using  $D^0 \rightarrow K^+ \pi^-$  Decays, *Phys. Rev. Lett.* **111**, 251801 (2013).
  - [11] R. Aaij *et al.* (LHCb Collaboration), First observation of  $D^0$ - $\bar{D}^0$  oscillations in  $D^0 \rightarrow K^+ \pi^+ \pi^- \pi^-$  decays and a measurement of the associated coherence parameters, *arXiv:1602.07224*.
  - [12] Y. Amhis *et al.* (Heavy Flavor Averaging Group), Averages of  $b$ -hadron,  $c$ -hadron, and  $\tau$ -lepton properties as of summer 2014, *arXiv:1412.7515*, updated results and plots available at <http://www.slac.stanford.edu/xorg/hfag/>.

- [13] T. Feldmann, S. Nandi, and A. Soni, Repercussions of flavor symmetry breaking on  $CP$  violation in  $D$ -meson decays, *J. High Energy Phys.* **06** (2012) 007.
- [14] B. Bhattacharya, M. Gronau, and J.L. Rosner,  $CP$  asymmetries in singly Cabibbo-suppressed  $D$  decays to two pseudoscalar mesons, *Phys. Rev. D* **85**, 054014 (2012).
- [15] D. Pirtskhalava and P. Uttayarat,  $CP$  violation and flavor  $SU(3)$  breaking in  $D$ -meson decays, *Phys. Lett. B* **712**, 81 (2012).
- [16] J. Brod, Y. Grossman, A.L. Kagan, and J. Zupan, A consistent picture for large penguins in  $D^0 \rightarrow \pi^- \pi^+$ ,  $K^- K^+$ , *J. High Energy Phys.* **10** (2012) 161.
- [17] H.-Y. Cheng and C.-W. Chiang,  $SU(3)$  symmetry breaking and  $CP$  violation in  $D \rightarrow PP$  decays, *Phys. Rev. D* **86**, 014014 (2012).
- [18] S. Müller, U. Nierste, and S. Schacht, Sum Rules of Charm  $CP$  Asymmetries beyond the  $SU(3)_F$  Limit, *Phys. Rev. Lett.* **115**, 251802 (2015).
- [19] M. Golden and B. Grinstein, Enhanced  $CP$  violations in hadronic charm decays, *Phys. Lett. B* **222**, 501 (1989).
- [20] H.-n. Li, C.-D. Lu, and F.-S. Yu, Branching ratios and direct  $CP$  asymmetries in  $D \rightarrow PP$  decays, *Phys. Rev. D* **86**, 036012 (2012).
- [21] R. Aaij *et al.* (LHCb Collaboration), Evidence for  $CP$  Violation in Time-Integrated  $D^0 \rightarrow h^- h^+$  Decay Rates, *Phys. Rev. Lett.* **108**, 111602 (2012).
- [22] Y. Grossman, A. L. Kagan, and Y. Nir, New physics and  $CP$  violation in singly Cabibbo suppressed  $D$  decays, *Phys. Rev. D* **75**, 036008 (2007).
- [23] M. Gronau,  $SU(3)$  in  $D$  decays: From 30% symmetry breaking to  $10^{-4}$  precision, *Phys. Rev. D* **91**, 076007 (2015).
- [24] Y. Grossman and D. J. Robinson,  $SU(3)$  sum rules for charm decay, *J. High Energy Phys.* **04** (2013) 067.
- [25] T. Aaltonen *et al.* (CDF Collaboration), Measurement of  $CP$ -violating asymmetries in  $D^0 \rightarrow \pi^+ \pi^-$  and  $D^0 \rightarrow K^+ K^-$  decays at CDF, *Phys. Rev. D* **85**, 012009 (2012).
- [26] M. Gersabeck, M. Alexander, S. Borghi, V. V. Gligorov, and C. Parkes, On the interplay of direct and indirect  $CP$  violation in the charm sector, *J. Phys. G* **39**, 045005 (2012).
- [27] A. L. Kagan and M. D. Sokoloff, Indirect  $CP$  violation and implications for  $D^0-\bar{D}^0$  and  $B_s^0-\bar{B}_s^0$  mixing, *Phys. Rev. D* **80**, 076008 (2009).
- [28] R. Aaij *et al.* (LHCb Collaboration), Measurement of  $CP$  asymmetry in  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  decays, *J. High Energy Phys.* **07** (2014) 041.
- [29] T. Aaltonen *et al.* (CDF Collaboration), Measurement of the Difference in  $CP$ -Violating Asymmetries in  $D^0 \rightarrow K^+ K^-$  and  $D^0 \rightarrow \pi^+ \pi^-$  Decays at CDF, *Phys. Rev. Lett.* **109**, 111801 (2012).
- [30] B. Aubert *et al.* (BABAR Collaboration), Search for  $CP$  Violation in the Decays  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$ , *Phys. Rev. Lett.* **100**, 061803 (2008).
- [31] M. Staric *et al.* (Belle Collaboration), Search for a  $CP$  asymmetry in Cabibbo-suppressed  $D^0$  decays, *Phys. Lett. B* **670**, 190 (2008).
- [32] B. R. Ko,  $CP$  violation and mixing in the charm sector at Belle, and current HFAG averages, [arXiv:1212.5320](https://arxiv.org/abs/1212.5320).
- [33] R. Aaij *et al.* (LHCb Collaboration), Measurement of the  $D^\pm$  production asymmetry in 7 TeV  $pp$  collisions, *Phys. Lett. B* **718**, 902 (2013).
- [34] R. Aaij *et al.* (LHCb Collaboration), Measurement of the  $D_s^+-D_s^-$  production asymmetry in 7 TeV  $pp$  collisions, *Phys. Lett. B* **713**, 186 (2012).
- [35] A. A. Alves Jr. *et al.* (LHCb Collaboration), The LHCb detector at the LHC, *J. Instrum.* **3**, S08005 (2008).
- [36] R. Aaij *et al.* (LHCb Collaboration), LHCb detector performance, *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
- [37] M. Adinolfi *et al.*, Performance of the LHCb RICH detector at the LHC, *Eur. Phys. J. C* **73**, 2431 (2013).
- [38] W. D. Hulsbergen, Decay chain fitting with a Kalman filter, *Nucl. Instrum. Methods Phys. Res., Sect. A* **552**, 566 (2005).
- [39] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.116.191601> for further plots and details.
- [40] N. L. Johnson, Systems of frequency curves generated by methods of translation, *Biometrika* **36**, 149 (1949).
- [41] R. Aaij *et al.* (LHCb Collaboration), Measurement of the Semileptonic  $CP$  Asymmetry in  $B^0-\bar{B}^0$  Mixing, *Phys. Rev. Lett.* **114**, 041601 (2015).
- [42] R. Aaij *et al.* (LHCb Collaboration), Measurement of the  $\bar{B}^0-B^0$  and  $\bar{B}_s^0-B_s^0$  production asymmetries in  $pp$  collisions at  $\sqrt{s} = 7$  TeV, *Phys. Lett. B* **739**, 218 (2014).
- [43] R. Aaij *et al.* (LHCb Collaboration), Study of the productions of  $\Lambda_b^0$  and  $\bar{B}^0$  hadrons in  $pp$  collisions and first measurement of the  $\Lambda_b^0 \rightarrow J/\psi p K^-$  branching fraction, *Chin. Phys. C* **40**, 011001 (2016).
- [44] K. A. Olive *et al.* (Particle Data Group), Review of particle physics, *Chin. Phys. C* **38**, 090001 (2014).
- [45] R. Aaij *et al.* (LHCb Collaboration), Measurements of Indirect  $CP$  Asymmetries in  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  Decays, *Phys. Rev. Lett.* **112**, 041801 (2014).
- [46] R. Aaij *et al.* (LHCb Collaboration), Measurement of indirect  $CP$  asymmetries in  $D^0 \rightarrow K^- K^+$  and  $D^0 \rightarrow \pi^- \pi^+$  decays, *J. High Energy Phys.* **04** (2015) 043.
- [47] R. Aaij *et al.* (LHCb Collaboration), Measurement of mixing and  $CP$  violation parameters in two-body charm decays, *J. High Energy Phys.* **04** (2012) 129.

---

R. Aaij,<sup>39</sup> C. Abellán Beteta,<sup>41</sup> B. Adeva,<sup>38</sup> M. Adinolfi,<sup>47</sup> A. Affolder,<sup>53</sup> Z. Ajaltouni,<sup>5</sup> S. Akar,<sup>6</sup> J. Albrecht,<sup>10</sup> F. Alessio,<sup>39</sup> M. Alexander,<sup>52</sup> S. Ali,<sup>42</sup> G. Alkhazov,<sup>31</sup> P. Alvarez Cartelle,<sup>54</sup> A. A. Alves Jr.,<sup>58</sup> S. Amato,<sup>2</sup> S. Amerio,<sup>23</sup> Y. Amhis,<sup>7</sup> L. An,<sup>3,40</sup> L. Anderlini,<sup>18</sup> G. Andreassi,<sup>40</sup> M. Andreotti,<sup>17,a</sup> J. E. Andrews,<sup>59</sup> R. B. Appleby,<sup>55</sup> O. Aquines Gutierrez,<sup>11</sup> F. Archilli,<sup>39</sup> P. d'Argent,<sup>12</sup> A. Artamonov,<sup>36</sup> M. Artuso,<sup>60</sup> E. Aslanides,<sup>6</sup> G. Auriemma,<sup>26,b</sup> M. Baalouch,<sup>5</sup> S. Bachmann,<sup>12</sup> J. J. Back,<sup>49</sup> A. Badalov,<sup>37</sup> C. Baesso,<sup>61</sup> W. Baldini,<sup>17,39</sup> R. J. Barlow,<sup>55</sup> C. Barschel,<sup>39</sup> S. Barsuk,<sup>7</sup> W. Barter,<sup>39</sup> V. Batozskaya,<sup>29</sup> V. Battista,<sup>40</sup> A. Bay,<sup>40</sup> L. Beaucourt,<sup>4</sup> J. Beddow,<sup>52</sup> F. Bedeschi,<sup>24</sup> I. Bediaga,<sup>1</sup> L. J. Bel,<sup>42</sup> V. Bellec,<sup>40</sup>



N. Belloli,<sup>21,c</sup> I. Belyaev,<sup>32</sup> E. Ben-Haim,<sup>8</sup> G. Bencivenni,<sup>19</sup> S. Benson,<sup>39</sup> J. Benton,<sup>47</sup> A. Berezhnoy,<sup>33</sup> R. Bernet,<sup>41</sup> A. Bertolin,<sup>23</sup> F. Betti,<sup>15</sup> M.-O. Bettler,<sup>39</sup> M. van Beuzekom,<sup>42</sup> S. Bifani,<sup>46</sup> P. Billoir,<sup>8</sup> T. Bird,<sup>55</sup> A. Birnkraut,<sup>10</sup> A. Bizzeti,<sup>18,d</sup> T. Blake,<sup>49</sup> F. Blanc,<sup>40</sup> J. Blouw,<sup>11</sup> S. Blusk,<sup>60</sup> V. Bocci,<sup>26</sup> A. Bondar,<sup>35</sup> N. Bondar,<sup>31,39</sup> W. Bonivento,<sup>16</sup> A. Borgheresi,<sup>21,c</sup> S. Borghi,<sup>55</sup> M. Borisyak,<sup>66</sup> M. Borsato,<sup>38</sup> T. J. V. Bowcock,<sup>53</sup> E. Bowen,<sup>41</sup> C. Bozzi,<sup>17,39</sup> S. Braun,<sup>12</sup> M. Britsch,<sup>12</sup> T. Britton,<sup>60</sup> J. Brodzicka,<sup>55</sup> N. H. Brook,<sup>47</sup> E. Buchanan,<sup>47</sup> C. Burr,<sup>55</sup> A. Bursche,<sup>41</sup> J. Buytaert,<sup>39</sup> S. Cadeddu,<sup>16</sup> R. Calabrese,<sup>17,a</sup> M. Calvi,<sup>21,c</sup> M. Calvo Gomez,<sup>37,e</sup> P. Campana,<sup>19</sup> D. Campora Perez,<sup>39</sup> L. Capriotti,<sup>55</sup> A. Carbone,<sup>15,f</sup> G. Carboni,<sup>25,g</sup> R. Cardinale,<sup>20,h</sup> A. Cardini,<sup>16</sup> P. Carniti,<sup>21,c</sup> L. Carson,<sup>51</sup> K. Carvalho Akiba,<sup>2</sup> G. Casse,<sup>53</sup> L. Cassina,<sup>21,c</sup> L. Castillo Garcia,<sup>40</sup> M. Cattaneo,<sup>39</sup> Ch. Cauet,<sup>10</sup> G. Cavallero,<sup>20</sup> R. Cenci,<sup>24,i</sup> M. Charles,<sup>8</sup> Ph. Charpentier,<sup>39</sup> M. Chefdeville,<sup>4</sup> S. Chen,<sup>55</sup> S.-F. Cheung,<sup>56</sup> N. Chiapolini,<sup>41</sup> M. Chrzasczcz,<sup>41,27</sup> X. Cid Vidal,<sup>39</sup> G. Ciezarek,<sup>42</sup> P. E. L. Clarke,<sup>51</sup> M. Clemencic,<sup>39</sup> H. V. Cliff,<sup>48</sup> J. Closier,<sup>39</sup> V. Coco,<sup>39</sup> J. Cogan,<sup>6</sup> E. Cogneras,<sup>5</sup> V. Cogoni,<sup>16,j</sup> L. Cojocariu,<sup>30</sup> G. Collazuol,<sup>23,k</sup> P. Collins,<sup>39</sup> A. Comerma-Montells,<sup>12</sup> A. Contu,<sup>39</sup> A. Cook,<sup>47</sup> M. Coombes,<sup>47</sup> S. Coquereau,<sup>8</sup> G. Corti,<sup>39</sup> M. Corvo,<sup>17,a</sup> B. Couturier,<sup>39</sup> G. A. Cowan,<sup>51</sup> D. C. Craik,<sup>51</sup> A. Crocombe,<sup>49</sup> M. Cruz Torres,<sup>61</sup> S. Cunliffe,<sup>54</sup> R. Currie,<sup>54</sup> C. D'Ambrosio,<sup>39</sup> E. Dall'Occo,<sup>42</sup> J. Dalseno,<sup>47</sup> P. N. Y. David,<sup>42</sup> A. Davis,<sup>58</sup> O. De Aguiar Francisco,<sup>2</sup> K. De Bruyn,<sup>6</sup> S. De Capua,<sup>55</sup> M. De Cian,<sup>12</sup> J. M. De Miranda,<sup>1</sup> L. De Paula,<sup>2</sup> P. De Simone,<sup>19</sup> C.-T. Dean,<sup>52</sup> D. Decamp,<sup>4</sup> M. Deckenhoff,<sup>10</sup> L. Del Buono,<sup>8</sup> N. Déleage,<sup>4</sup> M. Demmer,<sup>10</sup> D. Derkach,<sup>66</sup> O. Deschamps,<sup>5</sup> F. Dettori,<sup>39</sup> B. Dey,<sup>22</sup> A. Di Canto,<sup>39</sup> F. Di Ruscio,<sup>25</sup> H. Dijkstra,<sup>39</sup> S. Donleavy,<sup>53</sup> F. Dordei,<sup>39</sup> M. Dorigo,<sup>40</sup> A. Dosil Suárez,<sup>38</sup> A. Dovbnya,<sup>44</sup> K. Dreimanis,<sup>53</sup> L. Dufour,<sup>42</sup> G. Dujany,<sup>55</sup> K. Dungs,<sup>39</sup> P. Durante,<sup>39</sup> R. Dzhelyadin,<sup>36</sup> A. Dziurda,<sup>27</sup> A. Dzyuba,<sup>31</sup> S. Easo,<sup>50,39</sup> U. Egede,<sup>54</sup> V. Egorychev,<sup>32</sup> S. Eidelman,<sup>35</sup> S. Eisenhardt,<sup>51</sup> U. Eitschberger,<sup>10</sup> R. Ekelhof,<sup>10</sup> L. Eklund,<sup>52</sup> I. El Rifai,<sup>5</sup> Ch. Elsasser,<sup>41</sup> S. Ely,<sup>60</sup> S. Esen,<sup>12</sup> H. M. Evans,<sup>48</sup> T. Evans,<sup>56</sup> A. Falabella,<sup>15</sup> C. Färber,<sup>39</sup> N. Farley,<sup>46</sup> S. Farry,<sup>53</sup> R. Fay,<sup>53</sup> D. Fazzini,<sup>21,c</sup> D. Ferguson,<sup>51</sup> V. Fernandez Albor,<sup>38</sup> F. Ferrari,<sup>15</sup> F. Ferreira Rodrigues,<sup>1</sup> M. Ferro-Luzzi,<sup>39</sup> S. Filippov,<sup>34</sup> M. Fiore,<sup>17,39,a</sup> M. Fiorini,<sup>17,a</sup> M. Firlej,<sup>28</sup> C. Fitzpatrick,<sup>40</sup> T. Fiutowski,<sup>28</sup> F. Fleuret,<sup>71</sup> K. Fohl,<sup>39</sup> P. Fol,<sup>54</sup> M. Fontana,<sup>16</sup> F. Fontanelli,<sup>20,h</sup> D. C. Forshaw,<sup>60</sup> R. Forty,<sup>39</sup> M. Frank,<sup>39</sup> C. Frei,<sup>39</sup> M. Frosini,<sup>18</sup> J. Fu,<sup>22</sup> E. Furfaro,<sup>25,g</sup> A. Gallas Torreira,<sup>38</sup> D. Galli,<sup>15,f</sup> S. Gallorini,<sup>23</sup> S. Gambaetta,<sup>51</sup> M. Gandelman,<sup>2</sup> P. Gandini,<sup>56</sup> Y. Gao,<sup>3</sup> J. García Pardiñas,<sup>38</sup> J. Garra Tico,<sup>48</sup> L. Garrido,<sup>37</sup> D. Gascon,<sup>37</sup> C. Gaspar,<sup>39</sup> L. Gavardi,<sup>10</sup> G. Gazzoni,<sup>5</sup> D. Gerick,<sup>12</sup> E. Gersabeck,<sup>12</sup> M. Gersabeck,<sup>55</sup> T. Gershon,<sup>49</sup> Ph. Ghez,<sup>4</sup> S. Gianì,<sup>40</sup> V. Gibson,<sup>48</sup> O. G. Girard,<sup>40</sup> L. Giubega,<sup>30</sup> V. V. Gligorov,<sup>39</sup> C. Göbel,<sup>61</sup> D. Golubkov,<sup>32</sup> A. Golutvin,<sup>54,39</sup> A. Gomes,<sup>1,m</sup> C. Gotti,<sup>21,c</sup> M. Grabalosa Gándara,<sup>5</sup> R. Graciani Diaz,<sup>37</sup> L. A. Granado Cardoso,<sup>39</sup> E. Graugés,<sup>37</sup> E. Graverini,<sup>41</sup> G. Graziani,<sup>18</sup> A. Grecu,<sup>30</sup> P. Griffith,<sup>46</sup> L. Grillo,<sup>12</sup> O. Grünberg,<sup>64</sup> B. Gui,<sup>60</sup> E. Gushchin,<sup>34</sup> Yu. Guz,<sup>36,39</sup> T. Gys,<sup>39</sup> T. Hadavizadeh,<sup>56</sup> C. Hadjivasiliou,<sup>60</sup> G. Haefeli,<sup>40</sup> C. Haen,<sup>39</sup> S. C. Haines,<sup>48</sup> S. Hall,<sup>54</sup> B. Hamilton,<sup>59</sup> X. Han,<sup>12</sup> S. Hansmann-Menzemer,<sup>12</sup> N. Harnew,<sup>56</sup> S. T. Harnew,<sup>47</sup> J. Harrison,<sup>55</sup> J. He,<sup>39</sup> T. Head,<sup>40</sup> V. Heijne,<sup>42</sup> A. Heister,<sup>9</sup> K. Hennessy,<sup>53</sup> P. Henrard,<sup>5</sup> L. Henry,<sup>8</sup> J. A. Hernando Morata,<sup>38</sup> E. van Herwijnen,<sup>39</sup> M. Heß,<sup>64</sup> A. Hicheur,<sup>2</sup> D. Hill,<sup>56</sup> M. Hoballah,<sup>5</sup> C. Hombach,<sup>55</sup> W. Hulsbergen,<sup>42</sup> T. Humair,<sup>54</sup> M. Hushchyn,<sup>66</sup> N. Hussain,<sup>56</sup> D. Hutchcroft,<sup>53</sup> D. Hynds,<sup>52</sup> M. Idzik,<sup>28</sup> P. Ilten,<sup>57</sup> R. Jacobsson,<sup>39</sup> A. Jaeger,<sup>12</sup> J. Jalocha,<sup>56</sup> E. Jans,<sup>42</sup> A. Jawahery,<sup>59</sup> M. John,<sup>56</sup> D. Johnson,<sup>39</sup> C. R. Jones,<sup>48</sup> C. Joram,<sup>39</sup> B. Jost,<sup>39</sup> N. Jurik,<sup>60</sup> S. Kandybei,<sup>44</sup> W. Kalso,<sup>6</sup> M. Karacson,<sup>39</sup> T. M. Karbach,<sup>39</sup> S. Karodia,<sup>52</sup> M. Kecke,<sup>12</sup> M. Kelsey,<sup>39</sup> I. R. Kenyon,<sup>46</sup> M. Kenzie,<sup>39</sup> T. Ketel,<sup>43</sup> E. Khairullin,<sup>66</sup> B. Khanji,<sup>21,39,c</sup> C. Khurewathanakul,<sup>40</sup> T. Kim,<sup>9</sup> S. Klaver,<sup>55</sup> K. Klimaszewski,<sup>29</sup> O. Kochebina,<sup>7</sup> M. Kolpin,<sup>12</sup> I. Komarov,<sup>40</sup> R. F. Koopman,<sup>43</sup> P. Koppenburg,<sup>42,39</sup> M. Kozeiha,<sup>5</sup> L. Kravchuk,<sup>34</sup> K. Kreplin,<sup>12</sup> M. Kreps,<sup>49</sup> P. Krokovny,<sup>35</sup> F. Kruse,<sup>10</sup> W. Krzemien,<sup>29</sup> W. Kucewicz,<sup>27,n</sup> M. Kucharczyk,<sup>27</sup> V. Kudryavtsev,<sup>35</sup> A. K. Kuonen,<sup>40</sup> K. Kurek,<sup>29</sup> T. Kvaratskheliya,<sup>32</sup> D. Lacarrere,<sup>39</sup> G. Lafferty,<sup>55,39</sup> A. Lai,<sup>16</sup> D. Lambert,<sup>51</sup> G. Lanfranchi,<sup>19</sup> C. Langenbruch,<sup>49</sup> B. Langhans,<sup>39</sup> T. Latham,<sup>49</sup> C. Lazzeroni,<sup>46</sup> R. Le Gac,<sup>6</sup> J. van Leerdam,<sup>42</sup> J.-P. Lees,<sup>4</sup> R. Lefèvre,<sup>5</sup> A. Leflat,<sup>33,39</sup> J. Lefrançois,<sup>7</sup> E. Lemos Cid,<sup>38</sup> O. Leroy,<sup>6</sup> T. Lesiak,<sup>27</sup> B. Leverington,<sup>12</sup> Y. Li,<sup>7</sup> T. Likhomanenko,<sup>66,65</sup> M. Liles,<sup>53</sup> R. Lindner,<sup>39</sup> C. Linn,<sup>39</sup> F. Lionetto,<sup>41</sup> B. Liu,<sup>16</sup> X. Liu,<sup>3</sup> D. Loh,<sup>49</sup> I. Longstaff,<sup>52</sup> J. H. Lopes,<sup>2</sup> D. Lucchesi,<sup>23,k</sup> M. Lucio Martinez,<sup>38</sup> H. Luo,<sup>51</sup> A. Lupato,<sup>23</sup> E. Luppi,<sup>17,a</sup> O. Lupton,<sup>56</sup> A. Lusiani,<sup>24</sup> F. Machefert,<sup>7</sup> F. Maciuc,<sup>30</sup> O. Maev,<sup>31</sup> K. Maguire,<sup>55</sup> S. Malde,<sup>56</sup> A. Malinin,<sup>65</sup> G. Manca,<sup>7</sup> G. Mancinelli,<sup>6</sup> P. Manning,<sup>60</sup> A. Mapelli,<sup>39</sup> J. Maratas,<sup>5</sup> J. F. Marchand,<sup>4</sup> U. Marconi,<sup>15</sup> C. Marin Benito,<sup>37</sup> P. Marino,<sup>24,39,i</sup> J. Marks,<sup>12</sup> G. Martellotti,<sup>26</sup> M. Martin,<sup>6</sup> M. Martinelli,<sup>40</sup> D. Martinez Santos,<sup>38</sup> F. Martinez Vidal,<sup>67</sup> D. Martins Tostes,<sup>2</sup> L. M. Massacrier,<sup>7</sup> A. Massafferri,<sup>1</sup> R. Matev,<sup>39</sup> A. Mathad,<sup>49</sup> Z. Mathe,<sup>39</sup> C. Matteuzzi,<sup>21</sup> A. Mauri,<sup>41</sup> B. Maurin,<sup>40</sup> A. Mazurov,<sup>46</sup> M. McCann,<sup>54</sup> J. McCarthy,<sup>46</sup> A. McNab,<sup>55</sup> R. McNulty,<sup>13</sup> B. Meadows,<sup>58</sup> F. Meier,<sup>10</sup> M. Meissner,<sup>12</sup> D. Melnychuk,<sup>29</sup> M. Merk,<sup>42</sup> A. Merli,<sup>22,o</sup> E. Michielin,<sup>23</sup> D. A. Milanes,<sup>63</sup> M.-N. Minard,<sup>4</sup> D. S. Mitzel,<sup>12</sup> J. Molina Rodriguez,<sup>61</sup> I. A. Monroy,<sup>63</sup> S. Monteil,<sup>5</sup> M. Morandin,<sup>23</sup> P. Morawski,<sup>28</sup> A. Mordà,<sup>6</sup> M. J. Morello,<sup>24,i</sup> J. Moron,<sup>28</sup> A. B. Morris,<sup>51</sup>



R. Mountain,<sup>60</sup> F. Muheim,<sup>51</sup> D. Müller,<sup>55</sup> J. Müller,<sup>10</sup> K. Müller,<sup>41</sup> V. Müller,<sup>10</sup> M. Mussini,<sup>15</sup> B. Muster,<sup>40</sup> P. Naik,<sup>47</sup> T. Nakada,<sup>40</sup> R. Nandakumar,<sup>50</sup> A. Nandi,<sup>56</sup> I. Nasteva,<sup>2</sup> M. Needham,<sup>51</sup> N. Neri,<sup>22</sup> S. Neubert,<sup>12</sup> N. Neufeld,<sup>39</sup> M. Neuner,<sup>12</sup> A. D. Nguyen,<sup>40</sup> C. Nguyen-Mau,<sup>40,p</sup> V. Niess,<sup>5</sup> R. Niet,<sup>10</sup> N. Nikitin,<sup>33</sup> T. Nikodem,<sup>12</sup> A. Novoselov,<sup>36</sup> D. P. O'Hanlon,<sup>49</sup> A. Oblakowska-Mucha,<sup>28</sup> V. Obraztsov,<sup>36</sup> S. Ogilvy,<sup>52</sup> O. Okhrimenko,<sup>45</sup> R. Oldeman,<sup>16,48,j</sup> C. J. G. Onderwater,<sup>68</sup> B. Osorio Rodrigues,<sup>1</sup> J. M. Otalora Goicochea,<sup>2</sup> A. Otto,<sup>39</sup> P. Owen,<sup>54</sup> A. Oyanguren,<sup>67</sup> A. Palano,<sup>14,q</sup> F. Palombo,<sup>22,o</sup> M. Palutan,<sup>19</sup> J. Panman,<sup>39</sup> A. Papanestis,<sup>50</sup> M. Pappagallo,<sup>52</sup> L. L. Pappalardo,<sup>17,a</sup> C. Pappenheimer,<sup>58</sup> W. Parker,<sup>59</sup> C. Parkes,<sup>55</sup> G. Passaleva,<sup>18</sup> G. D. Patel,<sup>53</sup> M. Patel,<sup>54</sup> C. Patrignani,<sup>20,h</sup> A. Pearce,<sup>55,50</sup> A. Pellegrino,<sup>42</sup> G. Penso,<sup>26,r</sup> M. Pepe Altarelli,<sup>39</sup> S. Perazzini,<sup>15,f</sup> P. Perret,<sup>5</sup> L. Pescatore,<sup>46</sup> K. Petridis,<sup>47</sup> A. Petrolini,<sup>20,h</sup> M. Petrizzo,<sup>22</sup> E. Picatoste Olloqui,<sup>37</sup> B. Pietrzyk,<sup>4</sup> M. Pikies,<sup>27</sup> D. Pinci,<sup>26</sup> A. Pistone,<sup>20</sup> A. Piucci,<sup>12</sup> S. Playfer,<sup>51</sup> M. Plo Casasas,<sup>38</sup> T. Poikela,<sup>39</sup> F. Polci,<sup>8</sup> A. Poluektov,<sup>49,35</sup> I. Polyakov,<sup>32</sup> E. Polcarpo,<sup>2</sup> A. Popov,<sup>36</sup> D. Popov,<sup>11,39</sup> B. Popovici,<sup>30</sup> C. Potterat,<sup>2</sup> E. Price,<sup>47</sup> J. D. Price,<sup>53</sup> J. Prisciandaro,<sup>38</sup> A. Pritchard,<sup>53</sup> C. Prouve,<sup>47</sup> V. Pugatch,<sup>45</sup> A. Puig Navarro,<sup>40</sup> G. Punzi,<sup>24,s</sup> W. Qian,<sup>56</sup> R. Quagliani,<sup>7,47</sup> B. Rachwal,<sup>27</sup> J. H. Rademacker,<sup>47</sup> M. Rama,<sup>24</sup> M. Ramos Pernas,<sup>38</sup> M. S. Rangel,<sup>2</sup> I. Raniuk,<sup>44</sup> G. Raven,<sup>43</sup> F. Redi,<sup>54</sup> S. Reichert,<sup>55</sup> A. C. dos Reis,<sup>1</sup> V. Renaudin,<sup>7</sup> S. Ricciardi,<sup>50</sup> S. Richards,<sup>47</sup> M. Rihl,<sup>39</sup> K. Rinnert,<sup>53,39</sup> V. Rives Molina,<sup>37</sup> P. Robbe,<sup>7,39</sup> A. B. Rodrigues,<sup>1</sup> E. Rodrigues,<sup>55</sup> J. A. Rodriguez Lopez,<sup>63</sup> P. Rodriguez Perez,<sup>55</sup> S. Roiser,<sup>39</sup> V. Romanovsky,<sup>36</sup> A. Romero Vidal,<sup>38</sup> J. W. Ronayne,<sup>13</sup> M. Rotondo,<sup>23</sup> T. Ruf,<sup>39</sup> P. Ruiz Valls,<sup>67</sup> J. J. Saborido Silva,<sup>38</sup> N. Sagidova,<sup>31</sup> B. Saitta,<sup>16,j</sup> V. Salustino Guimaraes,<sup>2</sup> C. Sanchez Mayordomo,<sup>67</sup> B. Sanmartin Sedes,<sup>38</sup> R. Santacesaria,<sup>26</sup> C. Santamarina Rios,<sup>38</sup> M. Santimaria,<sup>19</sup> E. Santovetti,<sup>25,g</sup> A. Sarti,<sup>19,r</sup> C. Satriano,<sup>26,b</sup> A. Satta,<sup>25</sup> D. M. Saunders,<sup>47</sup> D. Savrina,<sup>32,33</sup> S. Schael,<sup>9</sup> M. Schiller,<sup>39</sup> H. Schindler,<sup>39</sup> M. Schlupp,<sup>10</sup> M. Schmelling,<sup>11</sup> T. Schmelzer,<sup>10</sup> B. Schmidt,<sup>39</sup> O. Schneider,<sup>40</sup> A. Schopper,<sup>39</sup> M. Schubiger,<sup>39</sup> M.-H. Schune,<sup>7</sup> R. Schwemmer,<sup>39</sup> B. Sciascia,<sup>19</sup> A. Sciubba,<sup>26,r</sup> A. Semennikov,<sup>32</sup> A. Sergi,<sup>46</sup> N. Serra,<sup>41</sup> J. Serrano,<sup>6</sup> L. Sestini,<sup>23</sup> P. Seyfert,<sup>21</sup> M. Shapkin,<sup>36</sup> I. Shapoval,<sup>17,44,a</sup> Y. Shcheglov,<sup>31</sup> T. Shears,<sup>53</sup> L. Shekhtman,<sup>35</sup> V. Shevchenko,<sup>65</sup> A. Shires,<sup>10</sup> B. G. Siddi,<sup>17</sup> R. Silva Coutinho,<sup>41</sup> L. Silva de Oliveira,<sup>2</sup> G. Simi,<sup>23,s</sup> M. Sirendi,<sup>48</sup> N. Skidmore,<sup>47</sup> T. Skwarnicki,<sup>60</sup> E. Smith,<sup>54</sup> I. T. Smith,<sup>51</sup> J. Smith,<sup>48</sup> M. Smith,<sup>55</sup> H. Snoek,<sup>42</sup> M. D. Sokoloff,<sup>58,39</sup> F. J. P. Soler,<sup>52</sup> F. Soomro,<sup>40</sup> D. Souza,<sup>47</sup> B. Souza De Paula,<sup>2</sup> B. Spaan,<sup>10</sup> P. Spradlin,<sup>52</sup> S. Sridharan,<sup>39</sup> F. Stagni,<sup>39</sup> M. Stahl,<sup>12</sup> S. Stahl,<sup>39</sup> S. Stefkova,<sup>54</sup> O. Steinkamp,<sup>41</sup> O. Stenyakin,<sup>36</sup> S. Stevenson,<sup>56</sup> S. Stoica,<sup>30</sup> S. Stone,<sup>60</sup> B. Storaci,<sup>41</sup> S. Stracka,<sup>24,i</sup> M. Straticiuc,<sup>30</sup> U. Straumann,<sup>41</sup> L. Sun,<sup>58</sup> W. Sutcliffe,<sup>54</sup> K. Swientek,<sup>28</sup> S. Swientek,<sup>10</sup> V. Syropoulos,<sup>43</sup> M. Szczekowski,<sup>29</sup> T. Szumlak,<sup>28</sup> S. T'Jampens,<sup>4</sup> A. Tayduganov,<sup>6</sup> T. Tekampe,<sup>10</sup> G. Tellarini,<sup>17,a</sup> F. Teubert,<sup>39</sup> C. Thomas,<sup>56</sup> E. Thomas,<sup>39</sup> J. van Tilburg,<sup>42</sup> V. Tisserand,<sup>4</sup> M. Tobin,<sup>40</sup> J. Todd,<sup>58</sup> S. Tolk,<sup>43</sup> L. Tomassetti,<sup>17,a</sup> D. Tonelli,<sup>39</sup> S. Topp-Joergensen,<sup>56</sup> E. Tournefier,<sup>4</sup> S. Tourneur,<sup>40</sup> K. Trabelsi,<sup>40</sup> M. Traill,<sup>52</sup> M. T. Tran,<sup>40</sup> M. Tresch,<sup>41</sup> A. Trisovic,<sup>39</sup> A. Tsaregorodtsev,<sup>6</sup> P. Tsopelas,<sup>42</sup> N. Tuning,<sup>42,39</sup> A. Ukleja,<sup>29</sup> A. Ustyuzhanin,<sup>66,65</sup> U. Uwer,<sup>12</sup> C. Vacca,<sup>16,39,j</sup> V. Vagnoni,<sup>15</sup> G. Valenti,<sup>15</sup> A. Vallier,<sup>7</sup> R. Vazquez Gomez,<sup>19</sup> P. Vazquez Regueiro,<sup>38</sup> C. Vázquez Sierra,<sup>38</sup> S. Vecchi,<sup>17</sup> M. van Veghel,<sup>43</sup> J. J. Velthuis,<sup>47</sup> M. Veltri,<sup>18,t</sup> G. Veneziano,<sup>40</sup> M. Vesterinen,<sup>12</sup> B. Viaud,<sup>7</sup> D. Vieira,<sup>2</sup> M. Vieites Diaz,<sup>38</sup> X. Vilasis-Cardona,<sup>37,e</sup> V. Volkov,<sup>33</sup> A. Vollhardt,<sup>41</sup> D. Voong,<sup>47</sup> A. Vorobyev,<sup>31</sup> V. Vorobyev,<sup>35</sup> C. Voß,<sup>64</sup> J. A. de Vries,<sup>42</sup> R. Waldi,<sup>64</sup> C. Wallace,<sup>49</sup> R. Wallace,<sup>13</sup> J. Walsh,<sup>24</sup> J. Wang,<sup>60</sup> D. R. Ward,<sup>48</sup> N. K. Watson,<sup>46</sup> D. Websdale,<sup>54</sup> A. Weiden,<sup>41</sup> M. Whitehead,<sup>39</sup> J. Wicht,<sup>49</sup> G. Wilkinson,<sup>56,39</sup> M. Wilkinson,<sup>60</sup> M. Williams,<sup>39</sup> M. P. Williams,<sup>46</sup> M. Williams,<sup>57</sup> T. Williams,<sup>46</sup> F. F. Wilson,<sup>50</sup> J. Wimberley,<sup>59</sup> J. Wishahi,<sup>10</sup> W. Wislicki,<sup>29</sup> M. Witek,<sup>27</sup> G. Wormser,<sup>7</sup> S. A. Wotton,<sup>48</sup> K. Wraight,<sup>52</sup> S. Wright,<sup>48</sup> K. Wyllie,<sup>39</sup> Y. Xie,<sup>62</sup> Z. Xu,<sup>40</sup> Z. Yang,<sup>3</sup> J. Yu,<sup>62</sup> X. Yuan,<sup>35</sup> O. Yushchenko,<sup>36</sup> M. Zangoli,<sup>15</sup> M. Zavertyaev,<sup>11,u</sup> L. Zhang,<sup>3</sup> Y. Zhang,<sup>3</sup> A. Zhelezov,<sup>12</sup> A. Zhokhov,<sup>32</sup> L. Zhong,<sup>3</sup> V. Zhukov,<sup>9</sup> and S. Zucchelli<sup>15</sup>

(LHCb Collaboration)

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China<sup>4</sup>LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France<sup>5</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France<sup>6</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France<sup>7</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France<sup>8</sup>LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France<sup>9</sup>I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany<sup>10</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

- <sup>11</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany
- <sup>12</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- <sup>13</sup>School of Physics, University College Dublin, Dublin, Ireland
- <sup>14</sup>Sezione INFN di Bari, Bari, Italy
- <sup>15</sup>Sezione INFN di Bologna, Bologna, Italy
- <sup>16</sup>Sezione INFN di Cagliari, Cagliari, Italy
- <sup>17</sup>Sezione INFN di Ferrara, Ferrara, Italy
- <sup>18</sup>Sezione INFN di Firenze, Firenze, Italy
- <sup>19</sup>Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy
- <sup>20</sup>Sezione INFN di Genova, Genova, Italy
- <sup>21</sup>Sezione INFN di Milano Bicocca, Milano, Italy
- <sup>22</sup>Sezione INFN di Milano, Milano, Italy
- <sup>23</sup>Sezione INFN di Padova, Padova, Italy
- <sup>24</sup>Sezione INFN di Pisa, Pisa, Italy
- <sup>25</sup>Sezione INFN di Roma Tor Vergata, Roma, Italy
- <sup>26</sup>Sezione INFN di Roma La Sapienza, Roma, Italy
- <sup>27</sup>Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland
- <sup>28</sup>AGH—University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland
- <sup>29</sup>National Center for Nuclear Research (NCBJ), Warsaw, Poland
- <sup>30</sup>Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania
- <sup>31</sup>Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia
- <sup>32</sup>Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia
- <sup>33</sup>Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia
- <sup>34</sup>Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia
- <sup>35</sup>Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia
- <sup>36</sup>Institute for High Energy Physics (IHEP), Protvino, Russia
- <sup>37</sup>Universitat de Barcelona, Barcelona, Spain
- <sup>38</sup>Universidad de Santiago de Compostela, Santiago de Compostela, Spain
- <sup>39</sup>European Organization for Nuclear Research (CERN), Geneva, Switzerland
- <sup>40</sup>Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
- <sup>41</sup>Physik-Institut, Universität Zürich, Zürich, Switzerland
- <sup>42</sup>Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands
- <sup>43</sup>Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands
- <sup>44</sup>NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine
- <sup>45</sup>Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine
- <sup>46</sup>University of Birmingham, Birmingham, United Kingdom
- <sup>47</sup>H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom
- <sup>48</sup>Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
- <sup>49</sup>Department of Physics, University of Warwick, Coventry, United Kingdom
- <sup>50</sup>STFC Rutherford Appleton Laboratory, Didcot, United Kingdom
- <sup>51</sup>School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
- <sup>52</sup>School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
- <sup>53</sup>Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom
- <sup>54</sup>Imperial College London, London, United Kingdom
- <sup>55</sup>School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom
- <sup>56</sup>Department of Physics, University of Oxford, Oxford, United Kingdom
- <sup>57</sup>Massachusetts Institute of Technology, Cambridge, MA, United States
- <sup>58</sup>University of Cincinnati, Cincinnati, OH, United States
- <sup>59</sup>University of Maryland, College Park, MD, United States
- <sup>60</sup>Syracuse University, Syracuse, NY, United States
- <sup>61</sup>Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil (associated with Institution Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil)
- <sup>62</sup>Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China (associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
- <sup>63</sup>Departamento de Física, Universidad Nacional de Colombia, Bogotá, Colombia (associated with Institution LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France)
- <sup>64</sup>Institut für Physik, Universität Rostock, Rostock, Germany (associated with Institution Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)
- <sup>65</sup>National Research Centre Kurchatov Institute, Moscow, Russia (associated with Institution Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)

- <sup>66</sup>*Yandex School of Data Analysis, Moscow, Russia (associated with Institution Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)*
- <sup>67</sup>*Instituto de Fisica Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain (associated with Institution Universitat de Barcelona, Barcelona, Spain)*
- <sup>68</sup>*Van Swinderen Institute, University of Groningen, Groningen, The Netherlands (associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands)*

<sup>a</sup>Also at Università di Ferrara, Ferrara, Italy.

<sup>b</sup>Also at Università della Basilicata, Potenza, Italy.

<sup>c</sup>Also at Università di Milano Bicocca, Milano, Italy.

<sup>d</sup>Also at Università di Modena e Reggio Emilia, Modena, Italy.

<sup>e</sup>Also at LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.

<sup>f</sup>Also at Università di Bologna, Bologna, Italy.

<sup>g</sup>Also at Università di Roma Tor Vergata, Roma, Italy.

<sup>h</sup>Also at Università di Genova, Genova, Italy.

<sup>i</sup>Also at Scuola Normale Superiore, Pisa, Italy.

<sup>j</sup>Also at Università di Cagliari, Cagliari, Italy.

<sup>k</sup>Also at Università di Padova, Padova, Italy.

<sup>l</sup>Also at Laboratoire Leprince-Ringuet, Palaiseau, France.

<sup>m</sup>Also at Universidade Federal do Triângulo Mineiro (UFMT), Uberaba-MG, Brazil.

<sup>n</sup>Also at AGH—University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

<sup>o</sup>Also at Università degli Studi di Milano, Milano, Italy.

<sup>p</sup>Also at Hanoi University of Science, Hanoi, Vietnam.

<sup>q</sup>Also at Università di Bari, Bari, Italy.

<sup>r</sup>Also at Università di Roma La Sapienza, Roma, Italy.

<sup>s</sup>Also at Università di Pisa, Pisa, Italy.

<sup>t</sup>Also at Università di Urbino, Urbino, Italy.

<sup>u</sup>Also at P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.